

Clinical Natural Language Processing and Audio

Tristan Naumann, Microsoft Research

CSC2541HS GUEST LECTURE

Using Clinical Text is Crucial

- Doctors use **text** to communicate patient state.
- Notes often contain the **most important** information



__date__ 4:07 AM
CHEST (PORTABLE AP) Clip # __num__
Reason: ETT tube placement, progression of pulmonary process
Admitting Diagnosis: **NON-HODGKIN LYMPHOMA**

__hospital__ MEDICAL CONDITION:
64 year old man s/p allo BMT for follicular lymphoma intubated now with
worsening respiratory status
REASON FOR THIS EXAMINATION:
ETT tube placement, progression of pulmonary process

FINAL REPORT
HISTORY: BMT for **lymphoma with respiratory status worsening.**

FINDINGS: In comparison with study of __date__, the tip of the endotracheal tube now measures approximately 3.2 cm above the carina. Central catheter and nasogastric tube remain in place. There is continued mild enlargement of the cardiac silhouette in a patient with low lung volumes. Indistinctness of engorged pulmonary vessels is consistent with elevated pulmonary venous pressure. The possibility of supervening consolidation cannot be excluded if there is appropriate clinical symptomatology.

ID	Description
6112	left vent drain
2734	right vent drain
1726	HIGH MIN VENT
1496	HIGH MIN. VENT.
1488	HIGH MIN. VENTIL.
1599	HIGH MINUTE VENT.

Clinical Text Presents Unique Challenges

CONTEXT
MATTERS

ACRONYMS



__date__ 4:07 AM
CHEST (PORTABLE AP) Clip # __num__
Reason: ETT tube placement, progression of pulmonary process
Admitting Diagnosis: **NON-HODGKIN LYMPHOMA**

__hospital__ MEDICAL CONDITION:
64 year old man s/p allo BMT for follicular lymphoma intubated now with
worsening respiratory status
REASON FOR THIS EXAMINATION:
ETT tube placement, progression of pulmonary process

FINAL REPORT
HISTORY: BMT for **lymphoma with respiratory status worsening.**

FINDINGS: In comparison with study of __date__, the tip of the endotracheal tube
now measures approximately 3.2 cm above the carina. Central catheter and
nasogastric tube remain in place. There is continued mild enlargement of the
cardiac silhouette in a patient with low lung volumes. Indistinctness of
engorged pulmonary vessels is consistent with elevated pulmonary venous
pressure. The possibility of supervening consolidation cannot be excluded if
there is appropriate clinical symptomatology.

ID	Description
6112	left vent drain
2734	right vent drain
1726	HIGH MIN VENT
1496	HIGH MIN. VENT.
1488	HIGH MIN. VENTIL.
1599	HIGH MINUTE VENT.

Clinical Audio Presents New Opportunities

- Automatic transcription of patient conversations and synthesis of notes.



Liz Hendricks, female, 48

*Liz is married to Ed. Has a daughter, Hannah, at the UW and a dog, Gerdy.
"I watch television shows about cooking. I take pictures of my food."*

CC: Patient is a 48-year-old woman with a new diagnosis of breast cancer

History of Present Illness

This is my husband, Ed. I came with her husband, Ed.

So three weeks ago, you had a concerning screening mammogram, sounds like.

Yeah. And just after that, a mammogram with an ultrasound.

Then I had a needle biopsy.

Followed by genetic testing. And what else?

And then you had a CT, of your chest, which is the one I saw.

Review of Systems

You mentioned that you have headaches sometimes.

Any trouble swallowing, pain when you swallow, anything like that?

Okay. And how about have you noticed that your breathing has changed at all, any ...

Do you ever get any chest pain or that heart racing feeling?

How about your belly? Any trouble with abdominal pain, diarrhea, constipation or ...

Any other symptoms like light headedness or trouble sleeping?

Social History

I work from home, live in Bellevue, I do marketing technology.

Doctor

Hi, Liz. I'm Dr. Hansen. Nice to meet you.

Patient

Hi.

Doctor

Thanks for coming in.

Patient

This is my husband, Ed.

Doctor

Hi, great to meet you, Ed. Thanks for coming in, being a support today.

So I'm the oncologist here at this office. There's a team of us.

And you've been referred by your PCP to talk about your situation with your breast mass and next steps to go from there.

Part of this visit will be for me to get to know your story a little bit.

I've read through your chart but I'd like to kind of hear from you.

And then we'll talk about the next steps, about moving forward.

I'm guessing this has been kind of a crazy, scary journey for you thus far.

Patient

Yeah, it hasn't been a normal three weeks for sure.

Doctor

Yeah, I'm guessing that's true.

Patient

Yeah.

Clinical Audio Presents New Opportunities

- Voice is another modality for assessing patient state:
 - Disorders directly related to voice (e.g. vocal hyperfunction).^{1,2}

Learning to detect vocal hyperfunction from
ambulatory neck-surface acceleration features:
Initial results for vocal fold nodules

Marzyeh Ghassemi, Jarrad H. Van Stan, Daryush D. Mehta, *Member, IEEE*,
Matías Zañartu, *Member, IEEE*, Harold A. Cheyne II, Robert E. Hillman, and John V. Guttag

Voice Disorder Identification by Using Machine Learning Techniques

LAURA VERDE¹, GIUSEPPE DE PIETRO², (*Member, IEEE*),
AND GIOVANNA SANNINO², (*Member, IEEE*)

¹Department of Engineering, Centro Direzionale di Napoli, Parthenope University of Naples, 80143 Naples, Italy

²Institute of High Performance Computing and Networking, 80131 Naples, Italy

Corresponding author: Giovanna Sannino (giovanna.sannino@icar.cnr.it)

- Disorders that manifest through voice (e.g. dementia).^{3,4}

Journal of Alzheimer's Disease 49 (2016) 407–422
DOI 10.3233/JAD-150520
IOS Press

407

Linguistic Features Identify Alzheimer's Disease in Narrative Speech

Kathleen C. Fraser^a, Jed A. Meltzer^b and Frank Rudzicz^{a,c,*}

^aDepartment of Computer Science, University of Toronto, Toronto, Canada

^bRotman Research Institute, Toronto, Canada

^cToronto Rehabilitation Institute-UHN, Toronto, Canada

Learning multiview embeddings for assessing dementia

Chloé Pou-Prom^{1,2,3}, Frank Rudzicz^{1,2,3}

¹ Toronto Rehabilitation Institute - UHN, Toronto, Canada

² Vector Institute, Toronto, Canada

³ Department of Computer Science, University of Toronto, Canada

{ chloe, frank }@cs.toronto.edu

Handling Associate Editor: Peter Garrard


[1] Ghassemi et al. Learning to detect vocal hyperfunction from ambulatory neck-surface acceleration features: Initial results for vocal fold nodules.

[2] Verde et al. Voice Disorder Identification by Using Machine Learning Techniques

[3] Fraser et al. Linguistic Features Identify Alzheimer's Disease in Narrative Speech.

[4] Pou-Prom et al. Learning multiview embeddings for assessing dementia.

Language is Part of A Larger Ecosystem



Outline

- **What is clinical language?**
- Common natural language processing (NLP) tasks.
- Resources for working with clinical language.
- Selected applications.

Clinical Text != Biomedical Text

- Biomedical text uses medical language in published literature. E.g.,



- Clinical text is collected by care staff and describes patients. E.g.,



Discharge Summary

Name: [**Last Name**],[**Known firstname**] Unit No: [**Numeric Identifier**]

Admission Date: [**__DATE__**] Discharge Date: [**__DATE__**]

Date of Birth: [**__DATE__**] Sex: F

Service: CARDIOTHORACIC

Allergies:

Ampicillin

Attending:[**First Name**]

Addendum:

Pt did not go to rehab on Percocet, but on Ultram instead.

Discharge Medications:

1. Aspirin 81 mg Tablet, Delayed Release (E.C.) Sig: One (1) Tablet, Delayed Release (E.C.) PO DAILY (Daily).

Disp:*30 Tablet, Delayed Release (E.C.)(s)* Refills:*2*

2. Docusate Sodium 100 mg Capsule Sig: One (1) Capsule PO BID (2 times a day).

Disp:*60 Capsule(s)* Refills:*2*

3. Amiodarone 200 mg Tablet Sig: Two (2) Tablet PO BID (2 times a day): 400mg twice a day for 7 days then decrease to 400mg daily for 7 days then decrease to 200mg daily until follow up with cardiologist.

Disp:*60 Tablet(s)* Refills:*2*

4. Pantoprazole 40 mg Tablet, Delayed Release (E.C.) Sig: One (1) Tablet, Delayed Release (E.C.) PO Q24H (every 24 hours).

Disp:*30 Tablet, Delayed Release (E.C.)(s)* Refills:*2*

Physician

TITLE:

Chief Complaint:

24 Hour Events:

EKG - At [**_DATE_**]

[**_DATE_**]: Started Metoprolol 25 TID. Around 8pm Pt tachy to 130s, dyspneic. O2 requirement increased from 2 to 4L. EKG obtained showed afib. Given 50 of metoprolol. Ordered 15 IV Dilt but held it as HR came down to 80s. CXR shows increased pulmonary edema. Pt urinating well so held off on **Diuretics**.

ECHO: The left atrium is dilated. There is mild symmetric left ventricular hypertrophy. The left ventricular cavity is moderately dilated. Overall left ventricular systolic function is moderately depressed (LVEF= 30-40 %) secondary to akinesis of the basal septum and

Infusions:

Heparin Sodium - 450 units/hour

Other ICU medications:

Other medications:

Changes to medical and family history:

Review of systems is unchanged from admission except as noted below

Vital signs

Hemodynamic monitoring

Fluid balance

24 hours
Since 12 AM

Tmax: 36.7

C (98

Tcurrent: 36.6

C (97.9

HR: 81 (63 - 114) bpm

Nursing

Sinus bradycardia. Long QTc interval. Low voltage in the limb leads.
No previous tracing available for comparison.

Normal sinus rhythm with atrio-ventricular conduction delay. Poor R wave progression in leads V1-V3 consistent with possible old anteroseptal myocardial infarction. Compared to the previous tracing of [**__DATE__**] the QRS voltage in the anterolateral leads is more prominent possibly related to lead placement.

Sinus rhythm
Nonspecific intraventricular conduction delay
Possible anterior infarct - age undetermined
Lateral T wave changes are nonspecific
Since previous tracing of [**__DATE__**], no significant change

Radiology

[**__DATE__**]

CTA CHEST W&W/O C&RECONS, NON-CORONARY

Reason: PLease rule out acute PE

Admitting Diagnosis: RESPIRATORY DISTRESS

Contrast: OMNIPAQUE Amt: 100

Clip # [**Clip Number (Radiology)**]

[**__HOSPITAL__**] MEDICAL CONDITION:

55 year old man with respiratory failure with history of PE, CPOD laryngeal edema.

REASON FOR THIS EXAMINATION:

PLease rule out acute PE

No contraindications for IV contrast


WET READ: EHAb [**First Name**] [**__DATE__**]

No pulmonary embolus detected (interval resolution since [**__DATE__**]). Moderate dependent atelectasis, right greater than left, in the setting of intubation. Thymic tissue noted, atypical for this age.


FINAL REPORT

HISTORY: History of pulmonary embolism, COPD, and laryngeal edema, now with respiratory failure. Evaluate for pulmonary embolism.

MIMIC-III Note Categories



MIMIC-III Note Lengths



Clinical Text Presents Unique Challenges

- **Data access:** often perceived as high risk due to difficulty of de-identification.
- **Copy-paste:** existing workflows encourage the repetition of existing text.
- **Quality variance:** some text well-written for communication, some not.
- **Partial structure:** sometimes generated or copied from structure (e.g vitals).
- **And the previous challenges with language...**

Additional Considerations for Audio

- **Disfluency:** non-trivial disfluencies in spoken language.

“I think you should um take you know aspirin.”

- **Utterance Segmentation:** imperfect speech turns complicate context.

“I’d like you to take albuterol for a week, also do you have an upcoming competition? I would like you to avoid vigorous exercise.

Outline

- What is clinical language?
- **Common natural language processing (NLP) tasks.**
- Resources for working with clinical language.
- Selected applications.

NLP Tasks

- Part of Speech
- Parsing
- Named Entity Recognition
- Normalization
- Negation
- Uncertainty
- Word Sense Disambiguation
- Relation Classification
- Summarization
 - Extractive
 - Abstractive

NLP Tasks

- Part of Speech
- Parsing
- **Named Entity Recognition**
- **Normalization**
- **Negation**
- **Uncertainty**
- Word Sense Disambiguation
- **Relation Classification**
- Summarization
 - Extractive
 - Abstractive

Named Entity Recognition

- Identify mentions of semantic types within text:
 - Find **spans** that correspond to an entity.
 - Classify the correct semantic type from the span.
- Context is important to resolve ambiguity, e.g.:
 - Ms Jane Doe has a history of MS
 - Mr Hutchinson diagnosed with Hutchinson

Normalization / Entity Linking

- Given an entity mention, assign canonical identifier (e.g., from ontology)
- We want each of these to have the same meaning:
 - Patient diagnosed with *RA* (C0003873)
 - Patient diagnosed with *Rheumatoid Arthritis* (C0003873)
 - Patient diagnosed with *atrophic arthritis* (C0003873)

Negation & Uncertainty

- **Negation:** entity mention is negated.
 - Patient denies *foot joint pain*.
 - foot joint pain, negated
C0458239, negated
- **Uncertainty:** entity mention not definitive.
 - Results suggestive of *colorectal cancer*.
 - colorectal cancer, probable
C1527249, probably

Relations

- **Relations:** high level semantic types relating more than one mention, e.g.:
 - *DegreeOf(modifier, disease/disorder)*
 - *LocationOf(anatomical site, sign/symptom)*
 - *Disrupts(anatomical site, disease/disorder)*
 - *Treats(drug, gene, mutation)*
- Helpful in forming higher level conceptual understanding.

Example of a Pipeline

An example of a sentence discovered by the sentence boundary detector:

Fx of obesity but no fx of coronary artery diseases.

Tokenizer output – 11 tokens found:

Fx of obesity but no fx of coronary artery diseases .

Normalizer output:

Fx of obesity but no fx of coronary artery disease .

Part-of-speech tagger output:

Fx of obesity but no fx of coronary artery diseases .
NN IN NN CC DT NN IN JJ NN NNS .

Shallow parser output:

Fx of obesity but no fx of coronary artery diseases .
NP PP \NP/ \NP/ PP \NP/

Named Entity Recognition – 5 Named Entities found:

Fx of obesity but no fx of coronary artery diseases .
obesity (type=diseases/disorders, UMLS CUI=C0028754, SNOMED-CT codes=308124008 and 5476005)
coronary artery diseases (type=diseases/disorders, CUI=C0010054, SNOMED-CT=8957000)
coronary artery (type=anatomy, CUI(s) and SNOMED-CT codes assigned)
artery (type=anatomy, CUI(s) and SNOMED-CT codes assigned)
diseases (type=diseases/disorders, CUI = C0010054)

Status and Negation attributes assigned to Named Entities:

Fx of obesity but no fx of coronary artery diseases .
obesity (status = family_history_of; negation = not_negated)
coronary artery diseases (status = family_history_of, negation = is_negated)

Figure 1 Example sentence processed through cTAKES components 'family history of obesity but no family history of coronary artery diseases.'
Fx, family history.

Outline

- What is clinical language?
- Common natural language processing (NLP) tasks.
- **Resources for working with clinical language.**
- Selected applications.

How can we represent language?

Medical Ontologies and Lexicons

- **SNOMED CT:** Codes, terms, synonyms, and definitions of clinical terms.
- **RxNorm:** Nomenclature of clinical drugs produced by NLM.
- **MeSH:** Medical Subject Headings (e.g., medical literature).
- **LOINC:** Logical Observation Identifiers Names and Codes (e.g. labs).
- **CPT:** Current Procedural Terminology maintained by AMA (e.g. billing).
- **ICD:** International Classification of Disease maintained by WHO (e.g. billing).

One System to Rule Them All

- **SNOMED CT:** Codes, terms, synonyms, and definitions of clinical terms.
- **RxNorm:** Nomenclature of clinical drugs produced by NLM
- **MeSH:** Medical Subject Headings (e.g., medical literature)
- **LOINC:** Logical Observation Identifiers Names and Codes (e.g. labs)
- **CPT:** Current Procedural Terminology maintained by AMA (e.g. billing).
- **ICD:** International Classification of Disease (e.g. billing)
- ...
- + 200 more sources



Unified Medical Language System (UMLS)

History of the UMLS

[Lindberg & al., *Methods*, 1993]
[Humphreys & al., *JAMIA*, 1998]

- Started at National Library of Medicine, 1986
- “Long-term R&D project”
- Complementary to IAIMS

(Integrated Academic Information Management Systems)



- «[...] the UMLS project is an effort to overcome two significant barriers to effective retrieval of machine-readable information.
- The first is the variety of ways the same concepts are expressed in different machine-readable sources and by different people.
 - The second is the distribution of useful information among many disparate databases and systems.»

The UMLS consists of

Metathesaurus

1 million+
biomedical
concepts
from over 100
sources

Semantic Network

135 broad
categories and
54 **relationships**
between
categories

SPECIALIST Lexicon & Tools

lexical
information and
programs for
**language
processing**

3 Knowledge Sources
used separately or together

Metathesaurus: clusters terms by meaning

- Synonymous terms clustered into a concept
- Preferred term is chosen
- Unique identifier (CUI) is assigned

Addison's disease	Metathesaurus	PN	
Addison's disease	SNOMED CT	PT	363732003
Addison's Disease	MedlinePlus	PT	T1233
Addison Disease	MeSH	PT	D000224
Bronzed disease	SNOMED Intl 1998	SY	DB-70620
Deficiency; corticorenal, primary	ICPC2-ICD10 Thesaurus	PT	MTHU021575
Primary Adrenal Insufficiency	MeSH	EN	D000224
Primary hypoadreanlism syndrome, Addison	MedDRA	LT	10036696

C0001403

Addison's disease

Semantic Network

- 135 Semantic Types
 - Broad subject categories (Clinical Drug, Virus)
 - Ex:
 - **Addison's Disease**
 - **Semantic Type: Disease or Syndrome**
- 54 Semantic Relationships
 - Links between categories (isa, causes, treats)
 - Ex:
 - Virus **causes** Disease or Syndrome
- Types + Relationships
 - Form the structure of the semantic network
 - Broadly categorize the biomedical domain

Concept

cluster of synonymous terms

Concept
C0001621

Term
adrenal disease gland
L0001621

S0011232 *Adrenal Gland Diseases*
S0011231 Adrenal Gland Disease
S0000441 Disease of adrenal gland
S0481705 Disease of adrenal gland, NOS
S0220090 Disease, adrenal gland
S0044801 Gland Disease, Adrenal

Term
adrenal disorder gland
unspecified
L0041793

S0860744 *Disorder of adrenal gland, unspecified*
S0217833 Unspecified disorder of adrenal glands

Term
adrenal disorder
L0161347

S0225481 *ADRENAL DISORDER*
S0627685 DISORDER ADRENAL (NOS)

Term
adrenal disorder gland
L0181041

S0632950 *Disorder of adrenal gland*
S0354509 Adrenal Gland Disorders

Term
L0162317

S0226798 *SURRENALE, MALADIES*

FRE

Is there data available?

i2b2: Informatics for Integrating Biology at the Bedside

- 2006 Deidentification and Smoking Challenge
- 2008 Obesity Challenge
- 2009 Medication Challenge
- 2010 Relations Challenge
- 2011 Coreference Challenge
- 2012 Temporal Relations Challenge
- 2014 De-identification and Heart Disease Risk Factors Challenge

- Challenge format: Datasets are **annotated!**¹

[1] <https://www.i2b2.org/>

SemEval & ShARe/CLEF

- 2014 SemEval Task 7: Analysis of Clinical Text¹
 - Entity, acronym, abbreviation recognition, mapping to CUIs
- 2015 SemEval Task 14: Analysis of Clinical Text²
 - Entity, acronym, abbreviation recognition, mapping to CUIs
- 2015 SemEval Task 6: Clinical TempEval³
 - Timespan, event, and temporal relation
- 2015: CLEF eHealth Evaluation Lab Task 1a: Clinical Speech Recognition⁴
 - Minimize word detection errors for Australian nursing shift changes


[1] <http://alt.qcri.org/semeval2014/task7/>

[2] <http://alt.qcri.org/semeval2015/task14/>

[3] <http://alt.qcri.org/semeval2015/task6/>

[4] <https://sites.google.com/site/clefehealth2015/task-1/task-1a>

Health NLP (hNLP) Center

[Home](#) [Data sets](#) [Tools](#) [Research +](#) [Join](#) [About +](#)

Health Natural Language Processing (hNLP) Center

The Health Natural Language Processing (hNLP) Center targets a key challenge to current hNLP research and health-related human language technology development: the lack of health-related language data.

The Center's primary activities are to:

1. Provide a repository and a data curation, distribution and management point for health-related language resources
2. Support sponsored research programs and health-related language-based technology evaluations
3. Engage in collaborations with US and foreign researchers, institutions and data centers
4. Host and participate in various workshops


The data consists of de-identified clinical notes from several institutions. We have paid special attention to the de-identification process which included a combination of automatic and manual redacting of information.

To obtain a data set, you must be a [member](#).


Layered Annotations

Some data sets contain layers of annotations. Click an image below to expand it.


Entity Recognition




Semantic Role Labelling








Properties and Relations



Temporal



 Boston Children's Hospital
United every child is well COLUMBIA UNIVERSITY
ON THE CITY OF NEW YORK University of Colorado
Boulder

Health Natural Language Processing Center.

MIMIC-III

- Over 2M notes for ~50K patients.
- Notes are de-identified, but otherwise unannotated.



Are tools available?

NLP Tools

- cTAKES¹ (clinical Text Analysis and Knowledge Extraction System)
 - Commonly regarded as the standard.
 - Flexible, but may require significant configuration.
- MetaMap²
 - Designed to identify UMLS concepts in text using knowledge-intensive approach.
- MetaMap Lite³
 - Less rigorous than MetaMap, but much faster.
- Sophia⁴ (v3NLP Framework)
 - Transform text into structured data for quality improvement, research, population health surveillance, and decision support. Scalable out of the box

[1] <http://ctakes.apache.org/>

[2] <https://metamap.nlm.nih.gov/>

[3] <https://metamap.nlm.nih.gov/MetaMapLite.shtml>

[4] <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5019303/>

NLP Tools

- CLAMP¹ (Clinical Language Annotation, Modeling and Processing Toolkit)
- MedEx²
- MedLEE (Medical Language Extraction and Encoding System)
- CliNER⁴ (Clinical Named Entity Recognition System)
- ...

[1] <https://clamp.uth.edu/>

[2] <http://www.vumc.org/cpm/cpm-blog/medex-tool-finding-medication-information>


[3] <http://www.medlingmap.org/taxonomy/term/80>

[4] <https://github.com/text-machine-lab/CliNER>

cTAKES

- Default pipeline is a good starting point for many projects.


The patient underwent a CT scan in April which did not reveal lesions in his liver.



cTAKES Components

- Sentence boundary
- Tokenization (rule-based)
- Morphologic normalization (NLM'S LVG)
- POS tagging
- Shallow parsing
- Named Entity Recognition
 - Dictionary mapping (lookup algorithm)
 - Machine learning
 - Types: Diseases/Disorders, Signs/Symptoms, Anatomical Sites, Procedures, Medications
- Negation and context identification (NegEx)
- Relation Extraction
- Clinical Element Model (CEM) normalization

Apache cTAKES Component Dependencies



Outline

- What is clinical language?
- Common natural language processing (NLP) tasks.
- Tools for working with clinical language.
- **Selected applications.**

Automated Trigger for Sepsis

Automated Trigger for Sepsis


- Horng et al.¹ use vitals and triage nursing notes to predict sepsis.

Which of these is likely to develop sepsis?

- pt with fever, chills, N/V since friday after eating what he thought was undercooked meat. Unable to hold po's down. Fevers to 103
- 89 yo f s/p esophageal hernia repair 3/09 w/ ?g-tube placement now w/ c/o's n&v. family reports pt's appetite is decreased, no BM x3d. generally not feeling well, had a bad day.
- from the scene fall of horse landed on r thigh deformity iv fluid 100 fentanyl/ morphine 4. no head or neck pain/

[1] Horng et al. "Creating an automated trigger for sepsis, clinical decision support at emergency department triage using machine learning." PLOS ONE, 2017

Text is much more predictive than structured data




Learning a representation for reasoning about a patient

- Observation: The goal of the triage note is to summarize a patient's state to provide maximal *context* in which to understand future data.
- Question: Can we learn the latent space directly from the triage text?
- Solution: Use a topic model called latent Dirichlet allocation (LDA) to identify underlying latent space.

Latent Dirichlet Allocation (LDA)

- Generative model where each document is a mixture of a small number of topics (inferred), and each word (observed) is attributable to one of those topics.



Underlying Topics Make Sense

- Their topic distributions appeal to clinical intuition.


Topic distributions	Less likely
facial numbness droop weakness sided speech slurred face...	↑
rabies bat vaccine exposure shot here for in room prophylaxis...	
shoulder pain rom arm decreased limited pulse injury ...	
etoh found admits unable ambulate trauma fs no on drinking...	
gait unsteady steady dizziness feet ha stable alert well oriented...	Infection
vaginal discharge bleeding vag d/c gyn itching pelvic foul...	↓
throat sore swallowing voice fevers ear difficulty st swallow...	
cellulitis swelling redness with lle rle leg and fevers l lower...	
pna cough on pneumonia with cxr dx recent levaquin r/o...	
	More likely

Summary: Notes Contain Important Observations of Uninstrumented Information


- Topic representations capture important information from clinical notes.
- Topic representations add predictive value over existing structured vitals.

Transfer Predictive Models Across EHRs

Transfer Predictive Models Across EHRs





Solution: Map Semantically Similar Items to Shared Vocabulary



Identify semantically equivalent concepts

Clinical Concepts Underlie Human-Readable Metadata



All: {C1328319, C0003086, C0445456, C0918012, C0205091}

Spanning: {C1328319, C0205091}

Longest: {C1328319}

Aggregate Representation


CUI: c1328319

ankle brachial index left


CUIs: c0003086 c0445456 c0918012 c0205091

(1 ...)

Aggregate Representation




Aggregate Representation



Aggregate Representation

	<i>CUI:</i> c1328319					
	ankle brachial index left					
<i>CUIs:</i>	c0003086	c0445456	c0918012	c0205091		
	(1			...)
		c0003086 ankle	c0445456 brachial	c0918012 index	c0205091 left	c1328319 ABI
<i>All:</i>	(1	1	1	1	1 ...)
<i>Spanning:</i>	(0	0	0	1	1 ...)
<i>Longest:</i>	(0	0	0	0	1 ...)

Aggregate Representation




Data & Experimental Setup

- Data and Representation

- Data from **2 MIMIC EHR systems** (CareVue and MetaVision).
- *Item IDs* encode charted observations as **bag-of-events**.
- Shared *Item IDs* from hospital EHR, and ICU-specific *Item IDs*.
- *Item IDs* mapped to UMLS CUIs.



- Model

- L2-regularized Logistic Regression, 5-fold cross-validation on training sets to determine best hyperparameters





Experiment 1: Does Mapping *Item ID Bag-of-Events* To *CUI Bag-of-Events* Perform Well Within Each EHR?

- Mapping *Item IDs* to CUIs does not degrade performance within single EHR.



Experiment 2: Does Mapping *Item ID Bag-of-Events* To *CUI Bag-of-Events* Perform Better Across EHRs?

- Mapping *Item IDs* to CUIs improves performance consistently across EHRs.



Summary: Text Enables Generalizability

- Application proposed an approach to automatically map semantically similar concepts from different databases to a common vocabulary.
- Application demonstrated the utility of this approach across an EHR transition on a set of prediction tasks.

Establishing the Availability of Information

Establishing the Availability of Information

- Complex models perform well in many domains, but results may be shallow.



- Are we leveraging the information that we should in unstructured text to predict outcomes?

Predicting Clinical Outcomes

- Target the prediction of common clinical outcomes in the ICU.

Task	Classes				
In-Hospital Mortality	Survived: 20,062	Expired: 924			
Diagnosis	Sepsis: 350	IC Hemorrhage: 295	Pneumonia: 483	CAD: 523	GI Bleed: 300
Admission Type	Urgent: 17,390	Elective: 3,596			
Length of Stay	Short (< 1.5 days): 6,722	Medium (1-5-3.5 days): 8,126	Long (> 3.5 days): 6,138		

Predicting Intermediate Information

- Establish the availability of important intermediate information to different classes of models.


Task	Classes		
Age	< 50 years: 4,565	50 - 80 years: 12,272	80+ years: 4,149
Gender	Male: 11,982	Female: 9,004	
Ethnicity	White: 14,974	Non-white: 3,282	

Evaluating Model Representations


- Form representations using:
 - Bag-of-words
 - Aggregate Word Embeddings
 - LSTM



Bag-of-words



Aggregate Word Embeddings



LSTM


Evaluating Model Representations

- Form representations using:
 - Bag-of-words
 - Aggregate Word Embeddings
 - LSTM


the dog is on the table

0	0	1	1	0	1	1	1
are	cat	dog	is	now	on	table	the

Bag-of-words



Aggregate Word Embeddings




LSTM

Evaluating Model Representations


- Form representations using:
 - Bag-of-words
 - Aggregate Word Embeddings
 - LSTM



Bag-of-words



Aggregate Word Embeddings




LSTM

Evaluating Model Representations


- Form representations using:
 - Bag-of-words
 - Aggregate Word Embeddings
 - LSTM



Bag-of-words





Aggregate Word Embeddings



LSTM

Experimental Setup



patient is a 64 y/o man → patient is a AGE_BETWEEN_SIXTY_AND_SEVENTY man

Representation Performance Varies By Task

- Simple models do surprisingly well on many tasks.

Binary AUCs

Task	In-Hospital Mortality	Admission Type	Gender	Ethnicity
BoW	.821	.883	.914	.619
Embeddings	.814	.873	.836	.580
LSTM	.777	.870	.837	.533

Representation Performance Varies By Task

- Simple models do surprisingly well on many tasks.

Binary AUCs

Task	In-Hospital Mortality	Admission Type	Gender	Ethnicity
BoW	.821	.883	.914	.619
Embeddings	.814	.873	.836	.580
LSTM	.777	.870	.837	.533

Representation Performance Varies By Task

- Simple models do surprisingly well on many tasks.

Binary AUCs

Task	In-Hospital Mortality	Admission Type	Gender	Ethnicity
BoW	.821	.883	.914	.619
Embeddings	.814	.873	.836	.580
LSTM	.777	.870	.837	.533

Representation Performance Varies By Task

- Simple models do surprisingly well on many tasks.

Binary AUCs

Task	In-Hospital Mortality	Admission Type	Gender	Ethnicity
BoW	.821	.883	.914	.619
Embeddings	.814	.873	.836	.580
LSTM	.777	.870	.837	.533

Multiclass Macro F1s

Task	Diagnosis	Length of Stay	Age
BoW	.828	.724	.635
Embeddings	.828	.730	.544
LSTM	.836	.758	.450

Representation Performance Varies By Task

- Simple models do surprisingly well on many tasks.

Binary AUCs

Task	In-Hospital Mortality	Admission Type	Gender	Ethnicity
BoW	.821	.883	.914	.619
Embeddings	.814	.873	.836	.580
LSTM	.777	.870	.837	.533


Multiclass Macro F1s

Task	Diagnosis	Length of Stay	Age
BoW	.828	.724	.635
Embeddings	.828	.730	.544
LSTM	.836	.758	.450

Word Vector Embeddings for Age Cluster Together, Confusing Prediction

- The age embedding is averaged across other signals from other embeddings.
- Obscures information that clinicians expect our models to know for relevance.

Task	Age
BoW	.635
Embeddings	.544
LSTM	.450



Summary: A Single Representation is Not Sufficient

- Simple representations have surprisingly good predictive power.
- Complex representations do not always capture basic information.
- Words that appear in similar contexts may still have different meanings.

Clinical Natural Language Processing and Audio

Tristan Naumann, Microsoft Research

tristan@microsoft.com